

4. (Amended) A formulation according to claim 3 wherein the oil phase is in a range from 0.2 to 40 percent (weight/vol)[, preferably, from 4 to 25 percent, more preferably, from 8 to 18 percent and, most preferably, from 8 to 10 percent] of the final formulation.

5. (Amended) A formulation according to [any of claims 1 to 4] claim 1 wherein the fat emulsion product employed for the preparation of said formulation contains from 2 to 40 percent, (weight/vol),[ preferably, from 5 to 25 percent, more preferably from 7 to 20 percent] of oil phase, from 0.2 to 5 percent, (weight/vol), [preferably, from 0.6 to 2 percent, more preferably, from 0.5 to 1.5 percent] of emulsifier and an additive in an amount suitable to control osmolarity[, preferably in a range from 1.5 to 5 percent (weight/vol), preferably from 2 to 3 percent].

6. (Amended) A formulation according to [any of claims 1 to 5] claim 1 wherein the oil phase contains long chain fatty acids in the form of triglycerides in the following proportions by weight:

linoleic acid	40-70%
oleic acid	15-30%
palmitic acid	5-15%
linoleic acid	3-12%
stearic acid	2-6%

7. (Amended) A formulation according to [any claims 1 to 6] claim 1 wherein the fat emulsion product employed for the preparation of said formulation comprises a composition selected from those reported in the following tables:

	Fat emulsion product 1	Fat emulsion product 2	Fat emulsion product 3
Soybean oil (w/vol)	10%	20%	5%

Safflower oil (w/vol)	--	--	5%
Egg yolk phospholipids (w/vol)	1.2%	1.2%	1.2%
Glycerol (w/vol)	2.25%	2.25%	2.5%
Fatty acids composition of vegetable oils (w/vol)			
Linoleic acid	50%	50%	65.8%
Oleic acid	26%	26%	17.7%
Palmitic acid	10%	10%	8.8%
Linolenic acid	9%	9%	4.2%
Stearic acid	3.5%	3.5%	3.4%
Osmolarity (mOsm/L)	260	268	276
Approximate pH	8	8	8
Fat particle size (μm)	0.5	0.5	0.4
Caloric value (cal/ml)	1.1	2.0	1.1
Size (ml)	50, 100	50, 100	25, 50
	250 or	250 or	100, 200
	500	500	Or 500
	Fat emulsion product 4	Fat emulsion product 5	Fat emulsion product 6
Soybean oil (w/vol)	10%	10%	20%
Safflower oil (w/vol)	10%	--	--
Egg yolk phospholipids (w/vol)	1.2%	1.2%	1.2%
Glycerol (w/vol)	2.5%	2.5%	2.5%
Fatty acids composition of vegetable oil (w/vol)			
Linoleic acid	65.8%	54.5%	54.5%
Oleic acid	17.7%	22.4%	22.4%
Palmitic acid	8.8%	10.5%	10.5%
Linolenic acid	4.2%	8.3%	8.3%
Stearic acid	3.4%	4.2%	4.2%

Osmolarity (mOsm/L)	258	284	292
Approximate pH	8.3	8.3	8.3
Fat particle size ( $\mu\text{m}$ )	0.4	0.4	0.4
Caloric value (cal/ml)	2.0	1.1	2.0
Size (ml)	25, 50	100, 200	200 or
	200 or	Or 500	550
	500		

and water for injection q.s. to 100%.

8. (Amended) A formulation according of claims 6 [or 7] wherein soybean oil and/or cottonseed oil and/or safflower oil are totally or partially replaced by a mixture of long chain fatty acids in the form of triglycerides to form a composition wherein said fatty acids are present in the respective proportions indicated in said claims and, optionally part of the above oils or long chain fatty acids is substituted by medium chain ( $\text{C}_6\text{-C}_{12}$ ) triglycerides.

9. (Amended) A formulation according to [any of claims 1 to 8] claim 1 wherein the concentration of the oil phase is between 4 and 25% (weight/vol)[, preferably, between 8 and 18%, more preferably between 8 and 10%] of the final formulation.

11. (Amended) A formulation according to [any one of claims 1 to 10] claim 1 wherein ramoplanin is present at a concentration between 1 and 20 mg/ml[, preferably from 1.5 to 15 mg/ml, most preferably, from about 3 to about 5 mg/ml].

12. (Amended) A formulation according to [any of claims 1 to 11] claim 1 wherein the pH of the final formulation is lower than 8[, preferably, lower than 7].

13. (Amended) A formulation according to [any of claims 1 to 10] claim 1 wherein the pH of the final formulation is between 4 and 6.5.

SHB 14  
14. (Amended) A formulation according to [any one of claims 1 to 13] claim 1 for treatment of infections caused by agents that are susceptible to ramoplanin or an antibiotic of the ramoplanin family.

15. (Amended) A formulation according to [any one of claims 1 to 13] claim 1 for the treatment of serious Gram positive infections - such as bacteremia, endocarditis or pneumonia.

65  
a1  
16. (Amended) A formulation according to [any one of claims 1 to 13] claim 1 for the treatment of severe infections caused by Gram positive drug-resistant or multiresistant microorganisms such as coagulase positive and negative staphylococci, penicillin-resistant streptococci or glycopeptide resistant enterococci.

17. (Amended) A formulation according to [any one of claims 1 to 16] claim 1 wherein ramoplanin factor A<sub>2</sub> is present in an amount of at least 75%.

18. (Amended) A pharmaceutical composition which consists of a ready to use dosage form or of a kit comprising separate packagings or containers containing ramoplanin or a member of ramoplanin family and the fat emulsion product for constitution of a formulation according to [any of claims 1 to 17] claim 1.--

Please add the following claims:

--20. The formulation according to claim 3, wherein said vegetal oil are selected from the group consisting of soybean oil, cottonseed oil, safflower oil and mixtures thereof.

a2  
21. The formulation according to claim 3, wherein said phospholipids are from an egg source.

22. The formulation according to claim 21, wherein said phospholipids are selected from the group consisting of egg lecithin, soybean lecithin, and mixtures thereof.

23. The formulation according to claim 3, wherein said osmotic agent is selected from the group consisting of sorbitol, glycerol, xylitol and mixtures thereof.

24. The formulation according to claim 4, wherein the oil phase is in the range of from 4 to 25 percent of the final formulation.

25. The formulation according to claim 4, wherein the oil phase is in the range of from 8 to 18 percent of the final formulation.

26. The formulation according to claim 4, wherein the oil phase is in the range of from 8 to 10 percent of the final formulation.

27. The formulation according to claim 5, wherein the fat emulsion product contains from 5 to 25 percent of the oil phase.

28. The formulation according to claim 5, wherein the fat emulsion product contains from 7 to 20 percent of the oil phase.

29. The formulation according to claim 5, wherein the fat emulsion product contains from 0.6 to 2 percent of the emulsifier.

30. The formulation according to claim 5, wherein the additive is in an amount from 1.5 to 5 percent.

31. The formulation according to claim 5, wherein the additive is in an amount from 2 to 3 percent.

32. The formulation according to claim 9, wherein the concentration of the oil phase is from 8 to 18 percent of the final formulation.

33. The formulation according to claim 11, wherein the ramoplanin is present in a concentration of from 1.5 to 15 mg/ml.

34. The formulation according to claim 11, wherein the ramoplanin is present in a concentration of from 3 to 5 mg/ml.